

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions and listings of claims in the application:

1. (Currently amended) An inhibitor of histone deacetylase represented by formula (1):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted;

X is selected from the group consisting of C=O, C=CH₂, CH(OH), CH(OR¹), C=N(OH), and C=N(OR¹), where R¹ is alkyl, aryl, aralkyl, or acyl;

Y¹ is a C₃-C₇ alkylene, wherein said alkylene may be optionally substituted, and wherein one or two carbon atoms in the ~~linear~~ alkylene chain connecting X and W may be replaced with O, NR³, or S(O)_n, where R³ is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, or caramoyl, and n is 0, 1, or 2, provided that the atoms in Y¹ that are attached to X and W are carbon atoms, and further provided that Y¹ does not comprise an ester or amide linkage in the linear chain connecting X and W; and

W is selected from the group consisting of -C(O)-CH₂-SR², -C(O)-NH-OM, -NH-C(O)-NH-Z, and -C(O)-NH-Z, where

R² is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted;

M is hydrogen or a pharmaceutically acceptable cation;


Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, anilinylmethyl, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, C₁-C₄ alkyl, or C₁-C₄ alkoxy;

provided that X is C=CH₂, CH(OR), C=N(OH), or C=N(OR¹) when W is -C(O)-NH-OM and Cy is unsubstituted phenyl, dimethylaminophenyl, or methoxyphenyl; and further provided that when W is -C(O)-CH₂-SR², the carbon atom in Y¹ that is attached to W is unsubstituted or is substituted with other than amino, acylamino, alkoxycarbonyl, or carbamoyl.

2. (Original) The inhibitor of claim 1, wherein Cy is C₆-C₁₀ aryl or is a radical of a heterocyclic moiety selected from the group consisting of thiophene, benzothiophene, furan, benzofuran, pyridine, quinoline, indole, isoquinoline, thiazole, morpholine, piperidine, and piperazine, any of which groups may be optionally substituted.
3. (Original) The inhibitor of claim 2, wherein the aryl or heterocyclic moiety is substituted by one or two substituents independently selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₆-C₁₀ aryl, heteroaryl, heterocyclyl, (C₆-C₁₀)ar(C₁-C₆)alkyl, halo, nitro, hydroxy, C₁-C₆ alkoxy, C₆-C₁₀ aryloxy, heteroaryloxy, C₁-C₆ alkoxycarbonyl, C₆-C₁₀ aryloxycarbonyl, heteroaryloxycarbonyl, carboxy, and amino.
4. (Previously Amended) The inhibitor of claim 1, wherein Cy has the formula -Cy¹-CY² or Cy¹-G-Cy², wherein Cy¹ and Cy² are independently C₃-C₆ cycloalkyl, C₆-C₁₀ aryl, or a radical of a heterocyclic moiety, which groups optionally may be substituted, and G or O, NR³ or S(O)_n, where R³ is hydrogen alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, or carbamoyl, and n is 0, 1, or 2.
5. (Cancelled)
6. (Original) The inhibitor of claim 1, where X is selected from the group consisting of CH(OR¹), C=N(OH), and C=N(OR¹), where R¹ is C₁-C₆ alkyl, C₆-C₁₀ aryl, or (C₆-C₁₀)ar(C₁-C₆)alkyl.
7. (Currently amended) The inhibitor of claim 1, wherein one to ~~about~~ three carbon atoms of the alkylene are independently substituted with halo, oxo, oximino, nitro, haloalkyl,

alkyl, aralkyl, alkoxy, aryloxy, alkoxycarbonyl, carboxy, hydroxyalkyl, acyl, acyloxy, or cyano.

8. (Original) The inhibitor of claim 1, wherein Y^1 comprises an all carbon linear chain connecting X and W.
9. (Original) The inhibitor of claim 8, wherein the linear chain connecting X and W comprises a dienyl moiety, wherein the dienyl moiety is attached to W.
10. (Currently Amended) The inhibitor of claim 9, wherein Y^1 is selected from the group consisting of



11. (Original) The inhibitor of claim 8, wherein Y^1 is $-(CH_2)_m$, where m is 5, 6, or 7.
12. (Original) The inhibitor of claim 1, wherein one carbon atom in the linear chain connecting X and W is replaced with O, NR^3 , or $S(O)_n$.
13. (Original) The inhibitor of claim 12, wherein Y^1 is $-(CH_2)-S(O)_n-(CH_2)_p$, where n is 0, 1, or 2, and p is 3, 4, or 5.
14. (Original) The inhibitor of claim 1, wherein W is $-C(O)-NH-OM$, M being selected from the group consisting of hydrogen, sodium, potassium, magnesium, and calcium.
15. (Original) The inhibitor of claim 1, wherein W is $-C(O)-NH-Z$ or $NH-C(O)-NH-Z$, Z being unsubstituted 2-anilinyl or unsubstituted 2-pyridyl.
16. (Original) The inhibitor of claim 1, wherein W is $-C(O)-CH_2-SR^2$, R^2 being selected from the group consisting of C_1-C_6 alkyl, C_6-C_{10} aryl, $(C_6-C_{10})ar(C_1-C_6)alkyl$, $(C_1-C_6 alkyl)Carbonyl$, $(C_6-C_{10} aryl)carbonyl$, and $((C_6-C_{10})ar(C_1-C_6)alkyl)carbonyl$, wherein the aryl portion of any such groups may be optionally substituted.

17. (Original) The inhibitor of claim 16, wherein R^2 is selected from the group consisting of methyl, phenyl, benzyl, benzoyl, and acetyl.
18. (Currently Amended) An inhibitor of histone deacetylase ~~represent~~ represented by formula (2):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted;

Y^2 is C_5 - C_7 alkylene, wherein said alkylene maybe optionally substituted, and wherein one or two carbon atoms in the ~~linear~~ alkylene chain connecting CY and W may be replaced with O, NR^3 , or $S(O)_n$, where R^3 is hydrogen, alkyl, aryl, arakyl, acyl, alkoxycarbonyl, or carbamoyl, and n is 0, 1, or 2, provided that Y^2 does not comprise an ester or amide linkage in the linear chain connecting Cy and W; and

W is selected from the group consisting of $-C(O)-CH_2-SR^2$, $-NH-C(O)-NH-Z$, and $-C(O)-NH-Z$, where

R^2 is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted; and

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, anilinyethyl, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, C_1 - C_4 alkyl, or C_1 - C_4 alkoxy;

provided that when W is $-C(O)-CH_2-SR^2$, the carbon atom in Y^2 that is attached to W is unsubstituted or is substituted with other than amino, acylamino, alkoxycarbonyl, or carbamoyl.

19. (Original) The inhibitor of claim 18, wherein Cy is C₆-C₁₀ aryl or is a radical of a heterocyclic moiety selected from the group consisting of thiophene, benzothiophene, furan, benzofuran, pyridine, uinoline, idole, isoquinoline, thiazole, morphonline, piperidine, piperazine, quinazolinone, benzotriazinone, phthalimide, and dioxobenzoisoquinoline, any of which groups may be optionally substituted.
20. (Original) The inhibitor of claim 18, wherein the aryl or heterocyclic moiety is substituted by one or two substituents independently selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ haloalkyl, C₆-C₁₀ aryl, heteroaryl, heterocyclyl, (C₆-C₁₀)ar(C₁-C₆)alkyl, halo, nitro, hydroxyl, C₁-C₆ alkoxy, C₆-C₁₀ aryloxy, heteroaryloxy, C₁-C₆ alkoxycarbonyl, C₆-C₁₀ aryloxycarbonyl, heteroaryloxycarbonyl, carboxy, and amino.
21. (Previously Amended) The inhibitor of claim 20, wherein Cy has the formula -Cy¹-Cy² or Cy¹-G-Cy², wherein Cy¹ and Cy² are independently C₃-C₆ cycloalkyl, C₆-C₁₀ aryl, or a radical of a heterocyclic moiety, which groups optionally may be substituted, and G is O, NR³, or S(O)_n, where R³ is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, or carbamoyl, and n is 0, 1, or 2.
22. (Cancelled)
23. (Original) The inhibitor of claim 18, wherein one to about four carbon atoms of the alkylene are independently substituted with halo, oxo, oximino, nitro, haloalkyl, alkyl, aralkyl, alkoxy, aryloxy, alkoxycarbonyl, carboxy, hydroxyalkyl, acyl, acyloxy, or cyano.
24. (Original) The inhibitor of claim 18, wherein one carbon atom in the linear chain connecting Cy and W is replaced with O, NR³, or S(O)_n.
25. (Original) The inhibitor of claim 19, wherein one carbon atom in the linear chain connecting Cy and W is replaced with NR³, where R³ is selected from the group consisting of C₁-C₆ alkyl, C₆-C₁₀ aryl, (C₆-C₁₀)ar(C₁-C₆)alkyl, (C₁-C₆alkyl) oxycarbonyl,

(C₆-C₁₀ aryl)oxycarbonyl, ((C₆-C₁₀)ar(C₁-C₆)alkyl)oxycarbonyl, (C₁-C₆ alkyl)carbonyl, (C₆-C₁₀ aryl)carbonyl, and ((C₆-C₁₀)ar(C₁-C₆)alkyl)carbonyl.

26. (Original) The inhibitor of claim 18, wherein one or two carbon atoms in the linear chain connecting Cy and W are replaced by O.
27. (Original) The inhibitor of claim 18, wherein W is -C(O)-NH-Z or -NH-C(O)-NH-Z, Z being unsubstituted 2-anilinyl or unsubstituted 2-pyridyl.
28. (Original) The inhibitor of claim 18, wherein W is -C(O)-CH₂-SR², R² being selected from the group consisting of C₁-C₆ alkyl, C₆-C₁₀ aryl, (C₆-C₁₀)ar(C₁-C₆)alkyl, (C₁-C₆ alkyl)carbonyl, (C₆-C₁₀ aryl)carbonyl, and ((C₆-C₁₀)ar(C₁-C₆)alkyl)carbonyl, wherein the aryl portion of any such groups may be optionally substituted.
29. (Original) The inhibitor of claim 28, wherein R² is selected from the group consisting of methyl, phenyl, benzyl, benzoyl, and acetyl.
30. (Original) An inhibitor of histone deacetylase represented by formula (3):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted, provided that Cy is other than dimethylaminoaphthyl when Y³ is -(CH₂)₃-;

Y³ is C₂-C₆ alkylene, wherein said alkylene may be optionally substituted with one or more substituents independently selected from the group consisting of halo, hydroxyl, oxo, nitro, haloalkyl, alkyl, aralkyl, alkoxy, aryloxy, carboxy, hydroxyalkyl, acyl, acyloxy, and cyano; and

W is selected from the group consisting of $-C(O)-CH_2-SR^2$, $-C(O)-NH-OM$, $-NH-C(O)-NH-Z$, and $-C(O)-NH-Z$, where

R^2 is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted;

M is hydrogen or a pharmaceutically acceptable cation; and

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, anilinylmethyl, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, C_1-C_4 , alkyl, or C_1-C_4 alkoxy;

provided that Z does not have the formula $-(C_5H_5N)-NHC(O)-Y^3-NH-S(O)_2-Cy$.

31. (Original) The inhibitor of claim 30, wherein Cy is C_6-C_{10} aryl or is a radical of a heterocyclic moiety selected from the group consisting of thiophene, benzothiophene, furan, benzofuran, pyridine, quinoline, indole, isoquinoline, thiazole, morpholine, piperidine, and piperazine, any of which groups may be optionally substituted.
32. (Original) The inhibitor of claim 31, wherein the aryl or heterocyclic moiety is substituted by one or two substituents independently selected from the group consisting of C_1-C_4 alkyl, C_1-C_4 haloalkyl, C_6-C_{10} aryl, heteroaryl, heterocyclyl, $(C_6-C_{10})ar(C_1-C_6)alkyl$, halo, nitro, hydroxyl, C_1-C_6 alkoxy, C_6-C_{10} aryloxy, heteroaryloxy, C_1-C_6 alkoxycarbonyl, C_6-C_{10} aryloxycarbonyl, heteroaryloxycarbonyl, carboxy, and amino.
33. (Previously Amended) The inhibitor of claim 30, wherein Cy has the formula $-Cy^1-Cy^2$ or Cy^1-G-Cy^2 , wherein Cy^1 and Cy^2 are independently C_3-C_6 cycloalkyl, C_6-C_{10} aryl, or a radical of a heterocyclic moiety, which groups optionally may be substituted, and G is O, NR^3 , or $S(O)_n$, where R^3 is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, or carbamoyl, and n is 0, 1, or 2.
34. (Cancelled)

35. (Original) The inhibitor of claim 30, wherein Y^3 is a C_2 - C_6 alkylene optionally substituted with one or two non-hydrogen substituents independently selected from the group consisting of halo, hydroxyl, oxo, nitro, (halo)₁₋₅(C_1 - C_3)alkyl, C_1 - C_6 alkyl, (C_6 - C_{10})ar(C_1 - C_6)alkyl, C_1 - C_6 alkoxy, C_6 - C_{10} arloxy, carboxy, hydroxy (C_1 - C_6)alkyl, C_1 - C_6 alkylcarbonyl, C_6 - C_{10} arylcarbonyl, C_1 - C_6 alkylcarbonylkloxy, C_6 - C_{10} arylcarbonyloxy, and cyano.
36. (Original) The inhibitor of claim 33, wherein Y^3 is an optionally substituted saturated C_4 - C_5 alkylene.
37. (Original) The inhibitor of claim 30, wherein W is $-C(O)-NH-OM$, M being selected from the group consisting of hydrogen, sodium, potassium, magnesium, and calcium.
38. (Original) The inhibitor of claim 30, wherein W is $-C(O)-NH-Z$ or $-NH-C(O)-NH-Z$, Z being unsubstituted 2-anilinyl or unsubstituted 2-pyridyl.
39. (Original) The inhibitor of claim 30, where W is $-C(O)-CH_2-SR^2$, R^2 being selected from the group consisting of C_1 - C_6 alkyl, C_6 - C_{10} aryl, (C_6 - C_{10})ar(C_1 - C_6)alkyl, (C_1 - C_6 alkyl)carbonyl, (C_6 - C_{10} aryl)carbonyl, and ((C_6 - C_{10})ar(C_1 - C_6)alkyl)carbonyl, wherein the aryl portion of any such groups may be optionally substituted.
40. (Cancelled)
41. (Cancelled)
42. (Currently Amended) A pharmaceutical composition comprising an inhibitor of histone deacetylase represented by formula (1):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted;

X is selected from the group consisting of C=O, C=CH₂, CH(OH), CH(OR¹), C=N(OH), and C=N(OR¹), where R¹ is alkyl, aryl, aralkyl, or acyl;

Y¹ is a C₃-C₇ alkylene, wherein said alkylene may be optionally substituted, and wherein one or two carbon atoms in the ~~linear~~ alkylene chain connecting X and W may be replaced with O, NR³, or S(O)_n, where R³ is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, or carbamoyl, and n is 0, 1, or 2, provided that the atoms in Y¹ that are attached to X and to W are carbon atoms, and further provided that Y¹ does not comprise an ester or amide linkage in the linear chain connecting X and W; and

W is selected from the group consisting of -C(O)-CH₂-SR², -C(O)-NH-OM, -NH-C(O)-NH-Z, and -C(O)-NH-Z, where R² is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted;

M is hydrogen or a pharmaceutically acceptable cation;

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, aniliny, methyl, or pyridylmethyl, any of which group optionally may be substituted with halo, hydroxyl, amino, nitro, C₁-C₄ alkyl, or C₁-C₄ alkoxy; and

a pharmaceutically acceptable carrier, excipient, or diluent; provided that X is C=CH₂, CH(OR¹), C=N(OH), or C=N(OR¹) when W is -C(O)-NH-OM and Cy is unsubstituted phenyl, dimethylaminophenyl, or methoxyphenyl; and

further provided that when W is -C(O)-CH₂-SR², the carbon atom in Y¹ that is attached to W is unsubstituted or is substituted with other than amino, acylamino, alkoxycarbonyl, or carbamoyl.

43. (Currently Amended) A pharmaceutical composition comprising an inhibitor of histone deacetylase represented by formula (2):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted;

Y^2 is $\text{C}_5\text{-C}_7$ alkylene, wherein said alkylene may be optionally substituted, and wherein one or two carbon atoms in the ~~linear~~ alkylene chain connecting Cy and W may be replaced with O, NR^3 , or S(O)_n , where R^3 is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxycarbonyl, or carbamoyl, and n is 0, 1, or 2, provided that Y^2 does not comprise an ester or amide linkage in the linear chain connecting Cy and W; and

W is selected from the group consisting of $-\text{C(O)}-\text{CH}_2\text{-SR}^2$, $-\text{NH-C(O)-NH-Z}$, and $-\text{C(O)-NH-Z}$, where R^2 is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted; and

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, anilinylmethyl, or ~~pyridylmethyl~~ pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, $\text{C}_1\text{-C}_4$ alkyl, or $\text{C}_1\text{-C}_4$ alkoxy; and a pharmaceutically acceptable carrier, excipient, or diluent;

provided that when W is $-\text{C(O)}-\text{CH}_2\text{-SR}^2$, the carbon atom in Y^2 that is attached to W is unsubstituted or is substituted with other than amino, acylamino, alkoxycarbonyl, or carbamoyl.

44. (Original) A pharmaceutical composition comprising an inhibitor of histone deacetylase represented by formula (3):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted, provided that Cy is other than dimethylaminonaphthyl when Y^3 is $-(CH_2)_3-$;

Y^3 is C_2-C_6 alkylene, wherein said alkylene may be optionally substituted with one or more substituents independently selected from the group consisting of halo, hydroxyl, oxo, nitro, haloalkyl, alkyl, aralkyl, alkoxy, aryloxy, carboxy, hydroxyalkyl, acyl, acyloxy, and cyano; and

W is selected from the group consisting of $-C(O)-CH_2-SR^2$, $-C(O)-NH-OM$, $-NH-C(O)-NH-Z$, and $-C(O)-NH-Z$, where

R^2 is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted;

M is hydrogen or a pharmaceutically acceptable cation; and

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, aniliny, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, C_1-C_4 alkyl, or C_1-C_4 alkoxy; and

a pharmaceutically acceptable carrier, excipient, or diluent.

provided that Z does not have the formula $-C_5H_3N)-NHC(O)-Y^3-NH-S(O)_2-Cy$.

45. (Cancelled)
46. (Cancelled)
47. (Currently Amended) A method of inhibiting histone deacetylase in a cell, comprising contacting a cell in which inhibition of histone deacetylase is desired with an inhibitor of histone deacetylase represented by formula (1):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted;

X is selected from the group consisting of C=O, C-CH₂, CH(OH), CH(OR¹), C=N(OH), and C=N(OR¹), where R¹ is alkyl, aryl, aralkyl, or acyl;

Y¹ is a C₃-C₇ alkylene, wherein said alkylene may be optionally substituted, and wherein one or two carbon atoms in the ~~linear~~ alkylene chain connecting X and W may be replaced with O, NR³, or S(O)_n, where R³ is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxy carbonyl, or carbamoyl, and n is 0, 1, or 2, provided that the atoms in Y¹ that are attached to X and W are carbon atoms, and further provided that Y¹ does not comprise an ester or amide linkage in the linear chain connecting X and W; and

W is selected from the group consisting of -C(O)-CH₂-SR²-C(O)-NH-OM, -NH-C(O)-NH-Z, and -C(O)-NH-Z, where

R² is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted;

M is hydrogen or a pharmaceutically acceptable cation;

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, anilinylmethyl, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, C₁-C₄ alkyl, or C₁-C₄ alkoxy;

provided that X is C=CH₂, CH(OR¹), C=N(OH), or C=N(OR¹) when W is -C(O)-NH-OM and Cy is unsubstituted phenyl dimethylaminophenyl, or methoxyphenyl; and

further provided that when W is $-\text{C}(\text{O})-\text{CH}_2-\text{SR}^2$, the carbon atom in Y^1 that is attached to W is unsubstituted or is substituted with other than amino, acylamino, alkoxycarbonyl, or carbamoyl.

48. (Currently Amended) A method of inhibiting histone deacetylase in a cell, comprising contacting a cell in which inhibition of histone deacetylase is desired with an inhibitor of histone deacetylase represented by formula (2):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted;

Y^2 is $\text{C}_5\text{-C}_7$ alkylene, wherein said alkylene may be optionally substituted, and wherein one or two carbon atoms in the ~~linear~~ alkylene chain connecting Cy and W may be replaced with O, NR^3 , or $\text{S}(\text{O})_n$, where R^3 is hydrogen, alkyl, aryl, aralkyl, acyl, alkoxybcarbonyl, or carbamoyl, and n is 0, 1, or 2, provided that Y^2 does not comprise an ester or amide linkage in the linear chain connecting Cy and W; and

W is selected from the group consisting of $-\text{C}(\text{O})-\text{CH}_2\text{SR}^2$, $-\text{NH}-\text{C}(\text{O})-\text{NH}-\text{Z}$, and $-\text{C}(\text{O})-\text{NH}-\text{Z}$, where

R^2 is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted; and

Z is selected from the group consisting of aniliny, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, aniliny, methyl, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, $\text{C}^1\text{-C}^4$ alkyl, or $\text{C}_1\text{-C}_4$ alkoxy;

provided that when W is $-\text{C}(\text{O})-\text{CH}_2-\text{SR}^2$, the carbon atom in Y^2 that is attached to W is unsubstituted or is substituted with other than amino, acylamino, alkoxycarbonyl, or carbamoyl.

49. (Original) A method of inhibiting histone deacetylase in a cell, comprising contacting a cell in which inhibition of histone deacetylase is desired with an inhibitor of histone deacetylase represented by formula (3):



wherein

Cy is cycloalkyl, aryl, or a radical of a heterocyclic moiety, any of which may be optionally substituted, provided that Cy is other than dimethylaminonaphthyl when Y^3 is $-(\text{CH}_2)_3-$;

Y^3 is $\text{C}_2\text{-C}_6$ alkylene, wherein said alkylene may be optionally substituted with one or more substituents independently selected from the group consisting of halo, hydroxyl, oxo, nitro, haloalkyl, alkyl, aralkyl, alkoxy, aryloxy, carboxy, hydroxyalkyl, acyl, acyloxy, and cyano; and

W is selected from the group consisting of $-\text{C}(\text{O})-\text{CH}_2-\text{SR}^2$, $-\text{C}(\text{O})-\text{NH}-\text{OM}$, $-\text{NH}-\text{C}(\text{O})-\text{NH}-\text{Z}$, and $\text{C}(\text{O})-\text{NH}-\text{Z}$, where



R^2 is alkyl, aryl, aralkyl, or acyl, wherein the aryl portion of any such groups may be optionally substituted;

M is hydrogen or a pharmaceutically acceptable cation; and



Z is selected from the group consisting of anilinyll, pyridyl, thiazolyl, hydroxyphenyl, thiadiazolyl, anilinyllmethyl, or pyridylmethyl, any of which groups optionally may be substituted with halo, hydroxyl, amino, nitro, $\text{C}_1\text{-C}_4$ alkyl, or C-C_4 alkoxy;

provided that Z does not have the formula $-(C_5H_3N)-NHC(O)-Y^3-NH-S(O)_2-Cy$.

50. (Original) A method of inhibiting histone deacetylase in a cell, comprising contacting a cell in which inhibition of histone deacetylase is desired with an inhibitor of histone deacetylase represented by one of formulae (4)-(5):



51. (Original) A method of inhibiting histone deacetylase in a cell, comprising contacting a cell in which inhibition of histone deacetylase is desired with an inhibitor of histone deacetylase represented by one of formulae (6)-(7):



Claims 52 - 56 (Cancelled)